

Vaccinations for your Child

Contact

Tomobe Health Center 0296-77-9145

Kasama Health Center 0296-72-7711

Iwama Health Center 0299-45-7888

	Type		Number of Shots	Ages Eligible for Public Funding
Infant (0 to 6 years)	Hib Note 1)	Initial	3 (27–56 days apart)	Over 2 months and under 5 years old
		Booster	1	
	Infant Pneumococcal Note 1)	Initial	3 (over 27 days apart)	Over 2 months and under 5 years old
		Booster	1	
	DPT-IPV • diphtheria • pertussis • tetanus • polio	Stage 1 Initial	3 (20–56 days apart)	Over 3 months and under 7 years 6 months old
		Stage 1 Booster	1	
	BCG			1
Measles/Rubella	Stage 1		1	Over 1 and under 2 years old

	(Combined vaccine) Note 2)	Stage 2			1	During the last year of kindergarten, etc. The one year period right before starting elem. school (from Apr 1st to Mar 31st)
	Chickenpox (Varicella)	Initial			1	Over 1 and under 3 years old
		Booster			1	
	Japanese encephalitis Note 3)	Stage 1 Initial	2	(6–28 days apart)		
Stage 1 Booster		4 years old	Roughly 1 year after the first vaccination		1	
School Age (elementary and up)	Japanese encephalitis Note 3)	Stage 2	9 years old		1	Over 9 and under 13 years old
	Combined diphtheria/tetanus (DT)	Stage 2	11 years old		1	Over 11 and under 13 years old
	Cervical cancer prevention		First year of junior high school		3	6th year elementary student to 1st year high school student